

# Hilft Beckenbodentraining gegen Inkontinenz in der Schwangerschaft?

Hebammenforum 2015; 16: 546-548

## References

- Hampel C et al.: Prevalence and natural history of female incontinence. *Eur Urol* 1997; 3:12.
- Thom DH, Rortveit G: Prevalence of Postpartum Urinary Incontinence: A Systematic Review. *J WOCN* 2011; 4: 25-26.
- Dolan LM et al.: Stress incontinence and pelvic floor neurophysiology 15 years after the first delivery. *BJOG* 2003; 12: 1107-1114.
- Viktrup L, Rortveit G, Lose G: Risk of stress urinary incontinence twelve years after the first pregnancy and delivery. *Obstet Gynecol* 2006; 2: 248-254.
- Anger JT et al.: Increasing costs of urinary incontinence among female Medicare beneficiaries. *J Urol* 2006; 1: 247-251.
- Ward-Smith P: Issues. The cost of urinary incontinence. *Urol Nurs* 2009; 3: 188.
- United Kingdom. National Institute for Health and Clinical Excellence (NICE): Urinary incontinence: the management of urinary incontinence in women [online]; 2006 [accessed: 2011 December 20th]. Herunterzuladen unter [www.nice.org.uk/nicemedia/live/10996/30281/30281.pdf](http://www.nice.org.uk/nicemedia/live/10996/30281/30281.pdf).
- Deutsche Kontinenz-Gesellschaft: Beckenbodentraining [online]; 2011 [accessed: 2011 December 20th]. Herunterzuladen unter: [www.kontinenz-gesellschaft.de/index.php?id=186](http://www.kontinenz-gesellschaft.de/index.php?id=186).
- Johanna Briggs Institute: The Joanna Briggs Institute Best Practice Information Sheet: The effectiveness of pelvic floor muscle exercises on urinary incontinence in women following childbirth. *Nurs Health Sci* 2011; 3: 378-381.
- Neumann PB, Grimmer KA, Deenadayalan Y: Pelvic floor muscle training and adjunctive therapies for the treatment of stress urinary incontinence in women: a systematic review. *BMC Womens Health* 2006; 6: 11.
- Mørkved S, Bø K: Effect of postpartum pelvic floor muscle training in prevention and treatment of urinary incontinence: a one-year follow up. *BJOG* 2000; 8: 1022-1028.
- Glazener CM, et al: Conservative management of persistent postnatal urinary and faecal incontinence: randomised controlled trial. *BMJ* 2001; 7313: 593-596.
- Chiarelli P, Cockburn J: Promoting urinary continence in women after delivery: randomised controlled trial. *BMJ* 2002; 7348: 1241
- Hay-Smith J et al.: Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database Syst Rev* 2008; 4: CD007471.
- Mason L et al.: The role of antenatal pelvic floor muscle exercises in prevention of postpartum stress incontinence: a randomised controlled trial. *J Clin Nurs* 2010; 19: 2777-2786.
- Dumoulin C, Hay-Smith EJC, Mac Habée-Séguin G: Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Cochrane Database Syst Rev* 2014. Report Nr Issue 5, Art. No.: CD005654.
- Boyle R, Hay-Smith EJC, Cody JD, Mørkved S: Pelvic floor muscle training for prevention and treatment of urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database Syst Rev* 2012. Report Nr Issue 10. Art. No.: CD007471.
- Stafne S et al.: Does regular exercise including pelvic floor muscle training prevent urinary and anal incontinence during pregnancy? A randomised controlled trial. *BJOG: An International Journal of Obstetrics and Gynaecology* 2012; 10: 1270-1280.
- Mørkved S et al.: Pelvic floor muscle training during pregnancy to prevent urinary incontinence: a single-blind randomized controlled trial. *Obstet Gynecol* 2003; 2: 313-319.
- Mørkved S, Bø K: The effect of postpartum pelvic floor muscle exercise in the prevention and treatment of urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 1997; 4: 217-222.
- Reilly ETC et al.: Prevention of postpartum stress incontinence in primigravidae with increased bladder neck mobility: a randomised controlled trial of antenatal pelvic floor exercises. *BJOG* 2002; 1: 68-76.
- Bø K, Haakstad LA: Is pelvic floor muscle training effective when taught in a general fitness class in pregnancy? A randomised controlled trial. *Physiotherapy* 2011; 3: 190-195.
- Farrar D, Tuffnell DJ, Ramage C: Interventions for women in subsequent pregnancies following obstetric anal sphincter injury to reduce the risk of recurrent injury and associated harms. *Cochrane Database Syst Rev* 2014. Report Nr Issue 11. Art. No.: CD010374.
- Mason L et al.: Do women practise pelvic floor exercises during pregnancy or following delivery? *Physiotherapy* 2001; 12: 662-670.
- Bø K, Hakstad L, Voldner N: Do pregnant women exercise their pelvic floor muscles? *Int Urogynecol J Pelvic Floor Dysfunct* 2007; 7: 733-736.
- Ismail SIMF: An audit of NICE guidelines on antenatal pelvic floor exercises. *Int Urogynecol J Pelvic Floor Dysfunct* 2009; 12: 1417-1422.
- Bø K: Pelvic floor muscle training is effective in treatment of female stress urinary incontinence, but how does it work. *International Urogynecology Journal* 2004; 76-84.
- Leonard K, Masatu MC: Outpatient process quality evaluation and the Hawthorne Effect. *Soc Sci Med* 2006; 9: 2330-2340.
- Huppelsberg J, Walter K: *Kurzlehrbuch Physiologie*. Thieme, 3. Aufl. 2009.
- Fine P et al.: Teaching and practicing of pelvic floor muscle exercises in primiparous women during pregnancy and the postpartum period. *Am J Obstet Gynecol* 2007; 1: 107.e1, 107.e5.
- Bump RC et al.: Assessment of Kegel pelvic muscle exercise performance after brief verbal instruction. *Am J Obstet Gynecol* 1991; 2: 322-327.
- Wilson PD, Herbison RM, Herbison GP: Obstetric practice and the prevalence of urinary incontinence three months after delivery. *Br J Obstet Gynaecol* 1996; 2: 154-161.
- Borello-France D et al.: Adherence to behavioral interventions for urge incontinence when combined with drug therapy: adherence rates, barriers, and predictors. *Phys Ther* 2010; 10: 1493-1505.